

# **Enhancing Hit Identification in *Mycobacterium tuberculosis* Drug Discovery Using Dual-Event Bayesian Models**

Sean Ekins<sup>1, 2\*</sup>, Robert C. Reynolds<sup>3,4</sup>, Scott G. Franzblau<sup>5</sup>, Baojie Wan<sup>5</sup> , Joel S. Freundlich<sup>6,7</sup> and Barry A. Bunin<sup>1</sup>

<sup>1</sup>Collaborative Drug Discovery, 1633 Bayshore Highway, Suite 342, Burlingame, CA 94010, USA.

<sup>2</sup>Collaborations in Chemistry, 5616 Hilltop Needmore Road, Fuquay-Varina, NC 27526, USA.

<sup>3</sup>Southern Research Institute, 2000 Ninth Avenue South, Birmingham, AL 35205, USA.

<sup>4</sup>Current address: University of Alabama at Birmingham, College of Arts and Sciences, Department of Chemistry, 1530 3<sup>rd</sup> Avenue South, Birmingham, Alabama 35294-1240, USA.

<sup>5</sup> Institute for Tuberculosis Research, University of Illinois at Chicago, Chicago, IL 60607, USA.

<sup>6</sup>Department of Medicine, Center for Emerging and Reemerging Pathogens, UMDNJ – New Jersey Medical School, 185 South Orange Avenue Newark, NJ 07103, USA.

<sup>7</sup>Department of Pharmacology & Physiology, UMDNJ – New Jersey Medical School, 185 South Orange Avenue Newark, NJ 07103, USA.

\*To whom correspondence should be addressed. (e-mail: [ekinssean@yahoo.com](mailto:ekinssean@yahoo.com))

**Running Head:** Dual Event Bayesian Models

**Table S1.** Mean (SD) leave one out and leave out 50% x 100 cross validation of previously published Bayesian models (ROC =receiver operator characteristic) – data from [22]

Dataset (number of molecules)	Leave one out ROC	Leave out 50% x 100 External ROC Score	Leave out 50% x 100Internal ROC Score	Leave out 50% x 100 Concordance	Leave out 50% x 100 Specificity	Leave out 50% x 100 Sensitivity
MLSMR						
All single point screen  (N = 220,463)	0.88	0.86 (0)	0.86 (0)	78.56 (1.86)	78.59(1.94)	77.13 (2.26)
MLSMR						
dose response set  (N = 2273)	0.78	0.73 (0.01)	0.75 (0.01)	66.85 (4.06)	67.21 (7.05)	65.47 (7.96)